

Suprep Colonoscopy Prep Instructions

Medications that must be stopped FIVE (5) days before the procedure:

Aspirin, Plavix (clopidogrel), Coumadin (warfarin), and Anti-Inflammatory drugs (eg: Motrin, Advil, Naproxen, Celebrex, Ibuprofen, Indocin, Aleve)

Medications that must be stopped THREE (3) days before the procedure: Xarelto (Rivaroxaban), Pradaxa (Dabigatran), Eliquis (Apixaban), Savaysa (Edoxaban), Brilinta (Ticagrelor)

Please check with your primary care physician or cardiologist whether you need Lovenox injections while you are off the blood thinner medications

Can be continued: (Tylenol, Percocet, Vicodin, Tramadol (Ultram), Oxycontin, Prednisone, Methadone, Codeine, Morphine, Neurontin, Fentanyl, Lidocaine are OKAY and can be continued.)

Day Before Colonoscopy

Refer to next page for diet plan

3 pm

Take first dose of Suprep

- Empty 1 bottle of liquid prep into 16oz mixing container that is supplied
- Add cold drinking water to the marked line - Mix to dissolve
- Drink - Take your time – finish within 1 hour
- Using a straw may help
- Drink 32 ounces of water over the next 3 hours

6 pm

Take 2 Dulcolax (bisacodyl) tablets

8 pm

Take 2 Dulcolax (bisacodyl) tablets

9 pm

Take second dose of Suprep

- Empty the 2nd bottle of liquid prep into the 16oz mixing container that is supplied
- Add cold drinking water to the marked line - Mix to dissolve
- Drink - Take your time – finish within 1 hours
- Using a straw may help
- Drink 32 ounces of water over the next 3 hours